

Norabent-F

Anti-caking agent/binding agent: 1m558i - Bentonite



▪ Properties

Appearence	White/pink powder
Classification	Technological additive for animal feed, classified in accordance with Implementing Regulation (EU) No 1060/2013 under additive code 1m558i, used as a binding agent and anti-caking agent.

▪ Chemical analysis

Smectite content	Min. 80 %
Moisture	8-12 %
Particule size > 75 µm	5-25 %
Swelling index	Min. 10ml/g
Cation exchange capacity (CEC), Ag-Th method	85 ± 10 meq /100g
pH	8,5 - 10

As - Arsenic	< 25 ppm
Hg - Mercury	< 1 ppm
Pb - Lead	< 30 ppm
Cd - Cadmium	< 2 ppm
F - Fluorine	< 4000 ppm
Dioxins (PCDD/F)	< 0.5 ng I-TEQ/kg
Dioxins (PCDD/F + Dioxin-like PCBs)	< 0.75 ng I-TEQ/kg
Non-dioxin PCBs	< 10 µg / kg

▪ Usage information

Directions for use	To be added to complete feeds for ruminants and poultry at a maximum inclusion rate of 20kg per ton.
Contra-indications and warnings	<p>The simultaneous use of orally administered macrolides should be avoided. For poultry, simultaneous use of robenidine should be avoided.</p> <ul style="list-style-type: none"> For poultry: the simultaneous use with coccidiostats other than robenidine is contraindicated with level of bentonite above 5000 mg/kg of complete feed. The total amount of bentonite must not exceed the permitted maximum level in complete feeding stuff of 20 000 mg/kg of complete feeding stuff. <p>Safety measures: When handling bentonite, it is recommended to wear appropriate respiratory protection, safety goggles and protective gloves to avoid inhalation of dust and any contact with the skin or eyes.</p>

▪ Other information

Packaging	25kg paper bag 1000/1250kg Big Bag
Storage	Store away from heat and humidity. Shelf life: 36 months after manufacture in original packaging.

α FR 56024003

