

Zinc Oxide 72 (Zno)



▪ Properties

Appearance	Brown powder
Classification	Nutritional additive trace element Zinc Oxide (3b603). Zinc (Zn) : 72 % min (typical value)

▪ Chemical analysis

Zn	72% min	Pb - Lead ⁽¹⁾	< 400 ppm
AS - Arsenic ⁽¹⁾	< 100,0 ppm	Dioxins ⁽¹⁾	< 1 ng/kg i-TEQ
Cd - Cadmium ⁽¹⁾	< 30,0 ppm	Dioxins and dioxin-like PCBs ⁽¹⁾	< 1,5 ng/kg i-TEQ
Hg - Mercury ⁽²⁾	< 0,5 ppm	Non-dioxin-like PCBs ⁽¹⁾	< 10 µg/kg i-TEQ

1) According to directive 2002/32/CE and amendments.

2) According to AFCA-CIAL recommendations.

▪ Usage information

Directions for use	Can be used in feed for all animal categories at a maximum content of Zinc (Zn) in mg/kg in the complete feed with 12% of moisture : <ul style="list-style-type: none"> • Dogs and cats: 200 (total) • Salmonids and milk replacers for calves: 180 (total) • Piglets, sows, rabbits and all fish other than salmonids: 150 (total) • Other species and categories: 120 (total). The additive shall be incorporated into feed in the form of a premixture.
Contra-indications and warnings	For users of the additive and premixtures, feed business operators shall establish operational procedures and appropriate organisational measures to address the potential risks by inhalation, dermal contact or eyes contact. Where risks cannot be reduced to an acceptable level by these procedures and measures, the additive and premixtures shall be used with appropriate personal protective equipment.

▪ Other information

Packaging	Bags of 25 kg Big Bag of 1000kg
Storage	Store away from heat sources and humidity. Shelf life : 36 months after manufacturing in original packaging

The information contained in this sheet is given to the best of our knowledge, without any guarantee on our part. Photo for illustrative purposes only.



α FR 56024003

